

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>A61K 39/12, C12N 7/08, A61K 35/76</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 98/33522</b> <b>(43) International Publication Date:</b> 6 August 1998 (06.08.98)
<b>(21) International Application Number:</b> PCT/EP98/00461 <b>(22) International Filing Date:</b> 29 January 1998 (29.01.98)  <b>(30) Priority Data:</b> 97200298.4                      3 February 1997 (03.02.97)                      EP <b>(34) Countries for which the regional or international application was filed:</b> NL et al.  <b>(71) Applicant (for all designated States except US):</b> DIMMINACO AG [CH/CH]; Zurichstrasse 12, CH-8134 Adliswil (CH).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> DAVELAAR, Frans, Gerrit [NL/NL]; Harderwijkstraat 85, NL-3881 EG Putten (NL). PITSTRA, Bernarda, Johanna [NL/NL]; Mouterspad 244, NL-1383 DK Weesp (NL). VAN WILTENBURG, Nico [NL/NL]; Koomanstraat 1G, NL-1391 VZ Abcoude (NL).  <b>(74) Agents:</b> MANNION, Sally, Kim; Wyeth Laboratories, Huntercombe Lane South, Taplow, Maidenhead, Berkshire SL6 0PH (GB) et al.	<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
<b>(54) Title:</b> INFECTIOUS BURSTITIS VACCINE  <b>(57) Abstract</b>  The present invention is concerned with a vaccine which is capable of protecting poultry against Infectious Bursitis Infections, characterized in that it contains an Infectious Bursitis virus which has the combined properties of upon administration to a chicken causing a reduction in the size of the bursal size, expressed as bursa/body weight ratio, of less than 55 % and the capability to protect poultry having an ELISA antibody titer of at least about 500 and with viruses having the above characteristics.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## INFECTIOUS BURSITIS VACCINE

5 The present invention is concerned with a vaccine which is capable of protecting poultry against Infectious Bursitis infections, and with novel viruses useful for preparing such vaccines.

10 Infectious Bursitis is an infective disease which can afflict poultry and which is caused by the Infectious Bursal Disease Virus (IBDV). The agent of Infectious Bursitis belongs to the family Birnaviridae, genus Birnavirus and affects primarily the Bursa of Fabricius. This causes an atrophy of the Bursa. Infection with the virulent viruses found in the field generally causes reduction of the bursa weight to approximately one-third of the normal weight. This is normally expressed as the bursa/body weight ratio (BBR). The disease gives degeneration and necrosis of the B lymphocyte population of the bursa. This may result in immunosuppression. The degree of atrophy (and subsequent degeneration) of  
15 the bursa depends on the virulence of the strain involved.

Serologically IBDV can be distinguished into two different types: type 1 is found in chickens, whereas type 2 is found in turkeys.

20 Prevention of Infectious Bursitis is based on vaccination. The vaccine strains currently in use are divided into highly virulent (= intermediate plus; examples: Bursine Plus, Bursine 3), intermediate virulent (e.g. Bursine 2, D78, LZ228TC) and highly attenuated mild (= avirulent; e.g. Bursine 1) strains.

25 The problem faced by poultry farmers is that in a regimen in which poultry flocks are regularly vaccinated, the newly hatched chicks are born with a high maternal antibody titre. It is known that maternal antibody titres interfere with the vaccine virus. Highly virulent, intermediate and avirulent strains break through maternal virus-neutralizing antibody levels of 1:500, 1:250 or less than 1:100, respectively ("Diseases of Poultry",  
30 ninth edition, ed: Calnek et al.; Iowa State University Press, 1991, p. 659). Using the ELISA system to monitor maternal antibodies, comparable results are obtained. The virulent (intermediate plus) vaccines break through IDDEX-ELISA titers of about 500. Intermediate and mild strains are found to break through much lower titers to induce a seroresponse.

35

The virulence of strains is monitored by the effect on the bursa weight (BBR) as described earlier, or by the microscopical changes. The following table (TABLE 1)

shows the level of virulence of intermediate plus vaccines (Poulvac Bursa Plus, Delvax LZ228E) and of virulent field virus (isolate D6948E) on the Bursa weight (expressed as percentage BBR as compared to normal).

TABLE 1

5

strain/vaccine	BBR 3 w. after inoculation (% of controls)
Poulvac Bursa Plus	27
Delvax LZ228E	39
D6948E 10a89	27
D6948E 10a89	39
mean	33

The comparison shows that the intermediate plus vaccines affect the bursa weight as much as the virulent field virus. As a consequence, practitioners who wish to vaccinate young chickens with high maternal antibody titers have to accept the negative  
10 consequences of the more "hot" strains (such as severe tracheal lesions and a high death rate) as well. Therefore, farmers lose chicks with > 1:500 antibody titre when using intermediate vaccines because it does not break through this titre, and when using a virulent vaccine, farms lose chicks because of the effect of the virus.

15 Hence, there is a need for a vaccine which is less virulent and capable of breaking through high maternal antibody titers at the same time.

Surprisingly, IBDV with this set of characteristics has been found. Thus, according to the present invention, there is provided a vaccine which is capable of protecting poultry  
20 against Infectious Bursitis infections, characterized in that it contains an Infectious Bursal Disease virus which has the combined properties of an intermediate virulence and the capability to protect poultry with an ELISA antibody titer of at least about 500.

A particular example of this new class of viruses has been isolated. The new IBDV  
25 strain is internally indicated as strain 9793, a sample of which has been deposited at the Collection Nationale de Cultures de Microorganismes of Institut Pasteur at Paris (France) under No. 1-1810 on January 22, 1997. The deposited material 1-1810 is in the form of the master seed virus.

30 The vaccine virus is prepared from Seed virus by one or more passages on a suitable medium. Preferably, all vaccine virus is derived from a single stock virus pool - the

Master Seed virus. Optionally a Working Seed Virus pool is prepared from this Master Seed by one or more passages, and the actual vaccine virus is prepared from this Working Seed by one or more passages as well. The vaccine normally contains a stabiliser, such as inositol or manitol, which is used in the preparation process. If  
5 required a diluent for the vaccine can be any physiological solution.

Suitable media for passaging during these respective propagation steps from Master Seed to vaccine virus are for example Specific Pathogen Free (SPF) eggs, Specific Antibody Negative (SAN) eggs, primary chicken cells or an avian cell line.

10 The vaccine according to the present invention preferably contains live viruses of the new IBDV class described above. The effective dose per chicken can vary between about  $10^2$  and about  $10^6$  EID<sub>50</sub>, preferably between about  $10^3$  and  $10^5$  EID<sub>50</sub>. The vaccine can be administered safely to one-day-old chicken and in ovo.

15 The vaccine according to the present invention based on live IBDV preferably is shipped in freeze-dried form and dissolved in water prior to use.

20 The vaccine according to the present invention preferably is administered by the enteral route, e.g. orally (in the drinking water or individually by pipette), oculonasally or by spray.

25 The IBDV strain 9793 is characterised by the ability to induce antibody in birds with a maternal antibody level of  $\geq 500$  (measured by IDDEX ELISA) and because it is much less damaging to the bursa than hot vaccines. Induction of antibody in birds with a maternal antibody level of  $\geq 500$  is normally only possible with so called "hot" vaccines and/or virulent field strains. These hot vaccines and field strains will badly damage the Bursa of Fabricius causing a depletion of B lymphocytes. IBDV strain 9793 is much less damaging to the bursa as measured by the size of the bursa after infection.

30 Thus, in further aspects of the invention there is provided a vaccine comprising a virus of the strain 9793, having the general or specific characteristics of deposit No. 1-1810, or a vaccine comprising a virus derived or developed from strain 9793, i.e. derived or developed from either the actual deposit or obtaining the virus from nature. There is also provided IBD virus having the general or specific characteristics of deposit No. 1-1810, or a vaccine comprising a virus derived or developed from strain 9793, i.e. derived or developed from either the actual deposit or obtaining the virus from nature.

The invention is illustrated, but not limited, by the following examples.

### EXAMPLE 1

#### Preparation of the master seed virus

5 Two vials of the mother material of strain 9793 were resuspended in 2.0 ml of distilled water. This virus suspension was diluted 1/10 by transferring 1 ml thereof into 9 ml of saline. After mixing this was diluted 1/30 by transferring 2 ml into 58 ml of saline to give a suspension containing approximately 500 EID<sub>50</sub> per 0.1 ml. Four hundred embryos of SAN eggs were each inoculated with 0.1 ml of the above virus suspension  
10 via the yolk sack route using a 1 ml syringe fitted with a 23 G x 1" needle. The eggs were incubated at 37°C and candled after 24 hours. Forty-nine embryos were dead and discarded. Forty-eight and seventy-two hours after inoculation the dead embryos (340 eggs total) were removed from the eggs and decapitated. They were pooled into groups of approximately 50 embryos. Each group was homogenized separately using an Ystral  
15 blender fitted with a small autoclavable shaft. Each homogenate was poured through two layers of gauze over a beaker and each filtrate was transferred to a separate Duran bottle. A volume of stabiliser, containing inositol and manitol, equivalent to approximately one third of the volume of the filtrate was added to each. Samples of 2 or 3 ml were taken from each bottle for titration or sterility testing respectively. The bottles  
20 and the samples were frozen and stored at -70°C until further processing. Sterile bottles were thawed, pooled and mixed well using a magnetic stirrer. A total volume of 496 ml of master seed suspension was filled into vials, 0.5 ml per vial, placed in the Edwards Lyomaster 4000 and freeze-dried. The vials were closed under vacuum with rubber stoppers and sealed with aluminium capsules.

25

### EXAMPLE 2

#### Preparation of the working seed.

One vial of the Master Seed Virus of strain 9793 was resuspended in 1.0 ml of distilled  
30 water. This was diluted 1/10 by transferring 0.2 ml thereof in 1.8 ml of saline. After mixing this was diluted 1/200 by transferring 1 ml in 199 ml of saline to give a suspension containing approximately 10<sup>4</sup> EID<sub>50</sub> per 0.1 ml. Six hundred and fifty embryos of SAN eggs were each inoculated with 0.1 ml of the above virus suspension via the yolk sack route using a 1 ml syringe fitted with a 23 G x 1" needle. The eggs  
35 were incubated at 37°C and candled after 24 and 48 hours. Twelve embryos were dead after 24 hours and eleven after 48 hours. These were all discarded. Seventy-two hours



after inoculation the embryos were removed from the remaining eggs and decapitated. They were pooled into ten groups of approximately 60 embryos. Each group was homogenized separately using an Ystral blender fitted with a small autocavable shaft. Each homogenate was poured through two layers of gauze over a beaker and each filtrate  
5 was transferred to a separate Duran bottle. A volume of stabiliser, containing inositol and manitol, equivalent to approximately one third of the volume of the filtrate was added to each. The final volume of each was between 50 and 60 ml. 1 ml was removed from each Duran bottle and pooled. After mixing this was dispensed in 1 ml aliquots into Nunc Cryotubes and these were stored at -70°C for later titration. A further 1 ml  
10 aliquot was removed from each Duran bottle and plated individually onto 5% Blood agar plates to check for sterility. The Duran bottles were all stored at -70°C until required. There was no growth on any of the plates after 14 days, therefore all ten aliquots were included in the working seed.

15 Freeze drying and testing of the Working Seed.

Each of the Working Seed aliquots were thawed, pooled and mixed well on a magnetic stirrer using a follower. Four hundred and twenty vials were each filled with 1 ml of virus suspension. The vials were placed into the Modulyo freeze drier. Freeze drying was carried out for fourteen hours before the vials were sealed and removed from the  
20 Clean Room. Crimp tops were applied to each vial and they were all tested for vacuum using an Edwards spark tester.

Preparation of the vaccine

The vaccine material itself was prepared according to the method as outlined above from  
25 the Working Seed. A stabiliser was added (inositol, manitol) to the vaccine material.

**EXAMPLE 3**

Safety of the vaccine

30 Safety of the vaccine prepared according to EXAMPLE 1 was tested after vaccination of one-day old chickens according to two methods explained below.

Bursa/body weight ratio

The bursa size is expressed as the bursa/body weight ratio (BBR) according to the  
35 following formula:

$$\text{BBR} = \frac{\text{bursa weight (g)} \times 1000}{\text{body weight (g)}}$$

Lesion score of bursa

The lesions seen by microscopical examination are judged as follows:

- 5 After weighing, the removed bursae were fixed in 4% formalin for microscopical examination at the PHC, Doorn, The Netherlands. The results were scored as summarized in TABLE 2.

TABLE 2

10

SCORE	OBSERVATIONS
0	No damage.
1	Some necrosis/lymphodepletion in isolated follicles.
2	Moderate generalized lymphodepletion or severe lymphodepletion in isolated follicles. Some fibrosis, some epithelial proliferation.
3	Over 50% of follicles with severe lymphodepletion. Fibrosis and epithelial proliferation. Some follicles have been replaced by cavities, initially filled with necrotic tissue (approx. 6 days post-infection), later on (after clearance of the necrotic tissue) lined with epithelium.
4	All follicles display severe lymphodepletion; only in their cortex lymphocytes may be discernible. Marked fibrosis and epithelial proliferation. Several follicles have been replaced by the afore-mentioned cavities.
5	Total loss of follicular architecture. Severe fibrosis and epithelial proliferation. The afore-mentioned cavities are scattered throughout the bursal tissue.

- 15 Groups of 30 birds each were vaccinated with the 9793 vaccine by spray, except for the controls. Three weeks after vaccination half of the birds of each group were killed and the BBR was determined. The remaining half of the birds of each group were challenged at 3 weeks after vaccination. This challenge was carried out using a highly virulent field strain (D6948E, obtained from the Animal Health Institute at Deventer, The Netherlands). 10-12 days after challenge these birds were sacrificed as well and the BBR was determined. From a selected number of groups both prior to and after challenge the bursa were examined microscopically and the lesion score was determined.



ResultsBBR.

The results of the BBR determinations are summarized in TABLE 3. These results show that the % BBR compared to the controls after vaccination as a mean is about 60.

- 5 Hence, it can be concluded that the damage to the bursa by the vaccine virus is very moderate. Furthermore, it is clear that the effect of the challenge virus to the bursa of vaccinated birds is absent, whereas in the control birds a considerable damage to the bursa is prominent. These latter data from the control birds in fact confirm the data earlier presented for field viruses and virulent vaccine strains.

10

TABLE 3

trial/ group	% BBR compared to the controls	
	3 w. after vaccination	10-12 days post-challenge
1a	46	47
1b	68	56
1c*	100	37
2a	45	55
2b	46	59
2c*	100	38
3a	51	62
3b	93	40
3c*	100	27
mean vaccinated	58.7	52
mean control	100	34.0

\* non-vaccinated controls.

Microscopical examination.

- 15 The microscopically visible damage to the bursa is summarized in TABLE 4. The lesion score after vaccination is very low and at an acceptable level for a vaccine, whereas a very high score is found in unvaccinated birds after challenge. The bursa of vaccinated birds again showed to be unaffected by the challenge virus.

TABLE 4

trial/ group	microscopical score	
	3 w. after vaccination	10 days post-challenge
3a	1.8	1.2
3b	1.5	1.2
3c*	0.6	4.2

\* non-vaccinated controls.

#### EXAMPLE 4

##### Seroresponse after vaccination

- 5 These studies were carried out with the birds used for EXAMPLE 3. Blood samples of the birds were taken at approximately 5 or 6 days of age. Individual titers 2-5 days later were calculated according to the formula of Kouwenhoven (B. Kouwenhoven and J. van den Bosch; in: Proceedings of 19th World's Poultry Congress, 1992, p. 465-468). Birds were divided then over different titer groups and vaccinated. During 3 or 4 weeks, blood
- 10 samples were taken to examine seroconversion.

#### Results

- The results of these studies are summarized in TABLE 5. These data confirm the high efficacy of the vaccine according to the present invention. High levels of maternal
- 15 antibodies do not interfere with the vaccination. Kouwenhoven (supra) gives ELISA titers of about 500 that intermediate plus strains are able to break through. The vaccine according to the present invention easily breaks through higher levels of maternal antibodies to induce seroconversion.

TABLE 5

trial group	calculated titer at day of vacc.	vacc. dose ( $10 \log \text{EID}_{50}$ )	mean antibody titer to IBD at days p.v.			
			7	14	21	28
1a	122	3.4	2	1061	1560	ND
1b	939	3.4	18	676	1088	ND
1c*	142	-	2	80	134	ND
2a	668	3.7	ND	3	410	1434
2b	649	3.7	ND	83	946	1808
2c*	1054	-	ND	85	3	31
3a	256	3.7	ND	760	2167	ND
3b	2235	3.7	ND	202	737	ND
3c*	1089	-	ND	42	55	ND

- 20 \* non-vaccinated controls; "ND" means not determined.

EXAMPLE 5

Bloodsamples were taken from one-day old chicks in order to calculate the vaccination day using the Kouwenhoven formula previously mentioned.

5 Two trials were set up, A and B, to determine effects of strain 9793 on mortality. For trial A bloodsamples were taken at one day old, at 14 (day of vaccination with strain 9793), 21, 28 and 35 days. Bloodsamples were taken at one day old, and at 11 (day of vaccination with strain 9793), 21, 28 and 35 days of age for trial B.

10 In both experiments one group was vaccinated for IBD and the other group acted as controls. Each group consisted of 1,500 chicks.

All chicks were vaccinated against Newcastle disease (ND), and Infectious Bronchitis (IB) at day of vaccination. Vaccines used were Poulvac NDW (Newcastle disease vaccine) and Poulvac IB Primer (Infectious Bronchitis vaccine containing strains H120 and D274). Vaccination was done by coarse spray. Serology was done by the HI  
15 test for ND and IB and by the IDEXX ELISA for IBD. Finally mortality was monitored. Data on serology are given in the Tables 6A and 6B (trial 97-62 A and 97-62 B).

Mortality was as follows:

20 Trial A: IBD vaccinated (group 1) = 7%  
unvacc. controls (group 2) = 6.4%

Trial B: IBD vaccinated (group 2) = 2.8%  
unvacc. controls (group 1) = 5.1%

25

Trial B shows that IBD 9793 breaks through maternal antibodies > 500. In trial A maternal antibodies were below 500 on the day of vaccination. Serology data for IB and ND do not show any immunosuppression by IBD 9793 on the humoral antibody response to IB or ND. Mortality data also demonstrate that the vaccine is safe.

30

Table 6A. Mean HI antibody titres to NDV, IBV M41 and IBV D 274 and mean ELISA antibody titres to IBDV.

age	no. sera	mean HI antibody titres			mean ELISA antibody titres
		ND	IB M 41	IB D274	IBD
on day					
group 1	24	4.2	7.7	7.3	3538
group 2	24	4.2	7.5	7.2	4208
14 days. Day of vaccination with stain 9793 (group 1)					
group 1	24	4.2	6.3	5.7	224
group 2	24	2.0	5.3	5.4	354
21 days. 1 week p.v.					
group 1	48	3.2	4.1	4.2	55
group 2	48	3.2	3.8	5.0	30
28 days. 2 weeks p.v.					
group 1	48	3.4	5.0	3.8	399 <sup>1</sup>
group 2	48	3.6	5.4	4.0	5
35 days. 3 weeks p.v.					
group 1	48	2.7	4.0	4.3	1708 <sup>1</sup>
group 2	48	3.2	4.2	4.6	4

1: statistically significant difference ( $P \leq 0.05$ ) between groups using one-way ANOVA with factor group.

Table 6B. Mean antibody titres against ND, IB M41, IB D274 and IBD

age	no. sera taken	mean HI antibody titres			mean ELISA antibody titres
		ND	IB M 41	IB D274	IBD
on day					
group 1	24	5.2	8.1	8.3	3113
group 2	24	4.9	8.5	8.7	3386
11 days. Day of vaccination with 9793 (group 2)					
<u>group 1</u>	<u>24</u>	<u>3.4</u>	<u>7.2</u>	<u>7.4</u>	<u>623</u>
<u>group 2</u>	<u>24</u>	<u>2.7</u>	<u>6.4</u>	<u>6.9</u>	<u>743</u>
21 days. 10 days p.v.					
group 1	48	3.1	5.4	5.7	25 <sup>1</sup>
group 2	48	3.8	5.2	5.8	157
28 days. 17 days p.v.					
group 1	48	3.3	4.2	6.6	35 <sup>1</sup>
group 2	48	3.2	4.1	6.4	773
35 days. 3 weeks p.v.					
group 1	48	3.6	3.9	5.4	89 <sup>1</sup>
group 2	48	4.0	4.0	5.6	1634

5

1: statistically significant difference ( $P \leq 0.05$ ) between groups using one-way ANOVA with factor group.

## CLAIMS

1. Vaccine which is capable of protecting poultry against Infectious Bursitis infections, characterized in that it comprises an Infectious Bursitis virus which has the combined properties of upon administration to a chicken of causing a reduction in bursal size, expressed as bursa/body weight ratio, of less than 55% and the capability to protect poultry having an ELISA antibody titer of at least about 500.
2. Vaccine comprising an Infectious Bursitis virus which has the characteristics of a virus of the strain 9793, a sample of which is deposited at Institut Pasteur at the Collection National de Cultures de Microorganismes under No. 1-1810.
3. Vaccine according to claim 1, characterized in that the Infectious Bursitis virus has the characteristics of deposit No. 1-1810.
4. Vaccine according to claim 1 or 2, characterized in that the Infectious Bursitis virus is a virus of the strain 9793, a sample of which is deposited at Institut Pasteur at the Collection National de Cultures de Microorganismes under No. 1-1810.
5. Vaccine according to any of claims 1 to 3, characterized in that the Infectious Bursitis virus is a virus derived or developed from strain 9793, a sample of which is deposited at Institut Pasteur at the Collection National de Cultures de Microorganismes under No. 1-1810.
6. Vaccine according to any of the preceding claims, characterized in that it contains the virus in an amount of between  $10^2$  and  $10^6$  EID<sub>50</sub> per dose.
7. Vaccine according to claim 6, characterized in that it contains the virus in an amount of between  $10^3$  and  $10^5$  EID<sub>50</sub> per dose.
8. Infectious Bursitis virus characterized by the combined properties of upon administration to a chicken causing a reduction in bursal size, expressed as bursa/body weight ratio, of less than 55% and the capability to protect poultry having an ELISA antibody titer of at least about 500.



9. Infectious Bursitis virus which has the characteristics of a virus of the strain 9793, a sample of which is deposited at Institut Pasteur at the Collection National de Cultures de Microorganismes under No. 1-1810.

10. Infectious Bursitis virus according to claim 8, characterized in that the Infectious Bursitis virus has the characteristics of deposit No. 1-1810.

11. Infectious Bursitis virus according to claim 8 or 9, characterized in that the Infectious Bursitis virus is a virus of the strain 9793, a sample of which is deposited at Institut Pasteur at the Collection National de Cultures de Microorganismes under No. 1-1810.

12. Infectious Bursitis virus according to any of claims 8 to 10, characterized in that the Infectious Bursitis virus is a virus derived or developed from strain 9793, a sample of which is deposited at Institut Pasteur at the Collection National de Cultures de Microorganismes under No. 1-1810.

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/EP 98/00461

**A. CLASSIFICATION OF SUBJECT MATTER**  
IPC 6 A61K39/12 C12N7/08 A61K35/76

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)  
IPC 6 A61K C12N C07K G01N C12P C12Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 600 723 A (ABIC LTD) 8 June 1994 see page 2, line 49 - page 3, line 44; claims 1.10	1,5
Y	---	1,3-5
X	US 5 192 539 A (VAN DER MAREL PIET ET AL) 9 March 1993 see claims 1-3	1,5
Y	---	1,3-5
X	US 4 824 668 A (MELCHIOR JR FRED W ET AL) 25 April 1989 see claims 2-24	1,5
Y	---	1,3-5
	--- -/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

**\* Special categories of cited documents :**

"A" document defining the general state of the art which is not considered to be of particular relevance  
"E" earlier document but published on or after the international filing date  
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)  
"O" document referring to an oral disclosure, use, exhibition or other means  
"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention  
"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone  
"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.  
"&" document member of the same patent family

Date of the actual completion of the international search

29 June 1998

Date of mailing of the international search report

09/07/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Halle, F

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 98/00461

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>LASHER H N ET AL: "Infectious bursal disease."</p> <p>WORLD'S POULTRY SCIENCE JOURNAL 50 (2). 1994. 133-166. ISSN: 0043-9339, XP002041802</p> <p>see page 149, paragraph Prevention of IBD</p> <p>---</p>	
A	<p>WO 91 05569 A (UNIV MARYLAND) 2 May 1991</p> <p>see claims 1-8</p> <p>-----</p>	

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 98/00461

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0600723 A	08-06-1994	IL 103939 A	12-09-1996
		BR 9304911 A	14-06-1994
		CN 1113158 A	13-12-1995
		CZ 9302605 A	15-06-1994
		HU 67324 A	28-03-1995
		PL 301253 A	13-06-1994
		ZA 9308952 A	02-08-1994
US 5192539 A	09-03-1993	AU 628063 B	10-09-1992
		AU 3825289 A	25-01-1990
		CA 1331444 A	16-08-1994
		CN 1039359 A	07-02-1990
		DE 68919449 D	05-01-1995
		DE 68919449 T	04-05-1995
		EP 0352835 A	31-01-1990
		ES 2067528 T	01-04-1995
		HU 211246 B	28-11-1995
		JP 2078634 A	19-03-1990
		JP 2749887 B	13-05-1998
		US 5602022 A	11-02-1997
US 4824668 A	25-04-1989	NONE	
WO 9105569 A	02-05-1991	AT 145236 T	15-11-1996
		AU 6605190 A	16-05-1991
		CA 2067397 A	19-04-1991
		DE 69029145 D	19-12-1996
		DE 69029145 T	07-05-1997
		EP 0496787 A	05-08-1992
		ES 2095258 T	16-02-1997
		JP 5504049 T	01-07-1993
		US 5518724 A	21-05-1996